

PHY-765 SS18 Gravitational Lensing Week 2

GL Geometry & Light Deflection

Kasper B. Schmidt

Leibniz-Institut für Astrophysik Potsdam (AIP)


Last week

- History of GL including early predictions including:
 - Light is affected by gravity
 - Deflections of 10s of arc sec for galaxy lenses
 - Useful for lens mass estimates
 - Spectroscopy is a key for identifying lenses
- People were considering deflection of light in Newtonian gravity (<1915):
 - $\alpha_N = 2GM / c^2 r$
- GR came along and changed this deflection to (>1915)
 - $\alpha_{GR} = 2 \times \alpha_N = 4GM / c^2 r$
- Growing importance of GL over the last 40 years (still growing)

The aim of today


- The Geometry of Gravitational Lensing:
 - Schematic of angles, distances and light-paths in GL
- Light deflection:
 - How did they arrive at the Newtonian deflection angle?
 - What changed things for the GR deflection angle?

Gravitational Lensing Geometry



The Newtonian Deflection Angle, α_N

- Last week we heard that, e.g., Newton, Cavendish, Laplace and Soldner all considered deflection of light in Newtonian gravity. But how?



- Consider a point particle, γ of mass m , deflected by a lens, L , with mass M
- Position a polar coordinate system (ϕ, R) with origin on the lens
- Let ϕ_0 be the angle at closest approach
- We know from Newtonian Gravity that in 3D

$$m \frac{d^2 \vec{r}}{dt^2} = - \frac{mMG}{r^2} \vec{r}$$

The Newtonian Deflection Angle, α_N

- But the motion considered here is confined to 2D (ϕ, R), so decomposing

$$m \frac{d^2 \vec{r}}{dt^2} = -\frac{mMG}{r^2} \vec{r}$$

- into the polar coordinate system just defined we have

$$\hat{R} \left[\ddot{R} - R\dot{\phi}^2 \right] + \cancel{\hat{\phi} \frac{1}{R} \frac{d}{dt} \left[R^2 \dot{\phi} \right]} = -\frac{MG}{R^2} \hat{R} \quad (\text{Exercise 2.1})$$

- Note: m drops out, i.e. this expression is independent of particle mass
 - Convenient given that we want to consider the mass-less photon, γ
- Angular momentum is a conserved quantity

$$R^2 \dot{\phi} = \frac{|L|}{m} \equiv J_z$$

- Hence

$$\frac{d}{dt} \left[R^2 \ddot{\phi} \right] = \frac{d}{dt} [J_z] = 0$$

The Newtonian Deflection Angle, α_N

- This gives

$$\begin{aligned}\hat{R} \left[\ddot{R} - R\dot{\phi}^2 \right] &= -\frac{MG}{R^2} \hat{R} \\ \ddot{R} - \frac{1}{R^3} \left(R^4 \dot{\phi}^2 \right) &= -\frac{MG}{R^2} \\ \ddot{R} - \frac{J_z^2}{R^3} &= -\frac{MG}{R^2}\end{aligned}$$

- Using that

$$\dot{R} = \frac{J_z R'}{R^2}$$

$$\ddot{R} = \frac{J_z^2}{R^2} \left[\frac{R''}{R^2} - 2 \frac{R'^2}{R^3} \right]$$

(Exercise 2.2)

- The equation of motion becomes

$$\frac{R''}{R^2} - 2 \frac{R'^2}{R^3} - \frac{1}{R} = -\frac{MG}{J_z^2}$$

The Newtonian Deflection Angle, α_N


- Changing variable from R to u , where $u \equiv 1/R$ we get

$$u'' + u = -\frac{MG}{J_z^2} \quad (\text{Exercise 2.3})$$

- This is an inhomogeneous second order differential equation
- The solution to this equation can be expressed by the cyclic function

$$\frac{1}{R(\phi)} = A \times \cos[\phi - \phi_0] + \frac{MG}{J_z^2}$$

- At very early times
 - R is large
 - ϕ is small



$$\tan \phi = \frac{\xi}{R} \simeq \phi \quad \Rightarrow \quad \frac{d}{dt} \phi = \dot{\phi} = -\frac{\xi}{R^2} \dot{R} \quad \Rightarrow$$

- Which means that the normalized angular momentum $J_z = -\xi \dot{R}$

The Newtonian Deflection Angle, α_N

- But as we are looking at the photon with velocity is $-c$ giving that

$$J_z = \xi c \quad (-c \text{ as velocity is opposite } R\text{-direction})$$

- which can be inserted into the solution to the equation of motion

$$\frac{1}{R(\phi)} = A \times \cos [\phi - \phi_0] + \frac{MG}{\xi^2 c^2}$$

- Still need to determine A and ϕ_0 . We will use “initial conditions” for this
- In the limit when $R \rightarrow \infty$ and $\phi \rightarrow 0$

$$\cos \phi_0 = \frac{MG}{A\xi^2 c^2}$$

The Newtonian Deflection Angle, α_N

- Secondly we look at the initial velocity, i.e. differentiating wrt. t

$$\frac{d}{dt} \frac{1}{R} = \frac{d}{dt} A \cos(\phi - \phi_0) + \frac{d}{dt} \frac{MG}{\xi^2 c^2}$$

\Downarrow

$$\begin{aligned} -\frac{\dot{R}}{R^2} &= A \left(-\sin(\phi - \phi_0) \dot{\phi} \right) \\ &= A \left(-\sin(\phi - \phi_0) \left(-\frac{\xi}{R^2} \dot{R} \right) \right) \end{aligned}$$

\Downarrow


$$A = \frac{1}{\sin(\phi - \phi_0)\xi} \rightarrow \frac{1}{\sin(\phi_0)\xi}$$

- This gives us two equations to determine the two unknowns A and ϕ_0

The Newtonian Deflection Angle, α_N

- So we have the following:

$$\frac{1}{R(\phi)} = A \times \cos[\phi - \phi_0] + \frac{MG}{\xi^2 c^2} \quad \cos \phi_0 = \frac{MG}{A \xi^2 c^2} \quad A = \frac{1}{\sin(\phi_0) \xi}$$



- What is the size of ϕ_0 without deflection?

$$\phi_0 = \pi/2$$

$$A \simeq \frac{1}{\xi} \Rightarrow \cos \phi_0 = \frac{MG}{\xi c^2}$$

- What is the size of the deflection compared to ϕ_0 ?

$$\phi \sim \pi/2 + \epsilon$$

- Taylor expanding this expression leads to:


$$\begin{aligned} \cos \phi_0 &\simeq \cos(\pi/2) + (-\sin(\pi/2))(\phi_0 - \pi/2) + \frac{-\cos(\pi/2)}{2}(\phi_0 - \pi/2)^2 + \dots \\ &\simeq \phi_0 - \pi/2 = -\epsilon \end{aligned}$$

The Newtonian Deflection Angle, α_N

- Which then results in

$$\phi_0 = \frac{\pi}{2} + \frac{MG}{\xi c^2}$$

- From geometry we can express α



$$2\psi + \alpha = \pi$$

$$(\pi - \phi_0) + \psi + \alpha = \pi$$

- So we have that

$$\alpha = 2\phi_0 - \pi$$

- Which combined with the above gives:

$$\alpha_N = \frac{2MG}{\xi c^2}$$

The GR Deflection Angle, α_{GR}

- Moving to GR, we want to describe the distortion by gravity in the curved space-time. Curvature analog to longitude and latitude on Earth

- Need to define some GR jargon:

- $g_{\mu\nu}$: The metric tensor where $g^{\mu\nu}$ is the inverse of $g_{\mu\nu}$
- $\Gamma^\beta_{\mu\nu}$: The affine connection, i.e. Christoffel symbols


$$\Gamma^\beta_{\mu\nu} = \frac{g^{\beta\beta}}{2} \left[\frac{\partial g_{\beta\mu}}{\partial x^\nu} + \frac{\partial g_{\beta\nu}}{\partial x^\mu} - \frac{\partial g_{\mu\nu}}{\partial x^\beta} \right]$$

- We consider the geodesic equation (geodesic = “straight line”)

$$\frac{d^2 x^i}{d\lambda^2} = -\Gamma^i_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}$$

- where $x^\alpha = (t, x, y, z)$ and β, μ, ν run over (0,1,2,3) and i, j, k over (1,2,3)
- In GR time t is not ‘special’ so differentiate wrt. the affine parameter λ

$$P = \frac{dx^\alpha}{d\lambda} \equiv p^\alpha = (E/c, \vec{p})$$



The GR Deflection Angle, α_{GR}

- Using that

$$\frac{dx^i}{d\lambda} = \frac{dx^i}{dt} \frac{dt}{d\lambda} = \frac{dx^i}{dt} \frac{E}{c}$$

$$\frac{d^2 x^i}{d\lambda^2} = \frac{E}{c} \frac{d}{dt} \left[\frac{E}{c} \frac{dx^i}{dt} \right] \simeq \frac{E^2}{c^2} \frac{d^2 x^i}{dt^2}$$
- We can write the geodesic equation as

$$\frac{E^2}{c^2} \frac{d^2 x^i}{dt^2} = -\Gamma_{\mu\nu}^i p^\mu p^\nu$$

- This is the expression for a particle's motion in a given space-time
- Need to define the space-time through the 'metric'.
- The metric when deflection is induced by a point mass M is

$$g_{\mu\nu} = \begin{pmatrix} g_{00} & 0 & 0 & 0 \\ 0 & g_{11} & 0 & 0 \\ 0 & 0 & g_{22} & 0 \\ 0 & 0 & 0 & g_{33} \end{pmatrix} \quad \text{where} \quad \begin{aligned} g_{00} &= c^2 \left(1 - \frac{2GM}{rc^2} \right) \\ g_{ij} &= -\delta_{ij} \left(1 + \frac{2GM}{rc^2} \right) \end{aligned}$$


The GR Deflection Angle, α_{GR}

- The line element for this metric is (analog to ‘Pythagoras’ in cartesian 2D)

$$ds^2 = g_{00} dt^2 + g_{ij} dx^i dx^j = 0$$

- GR time dilation:

$$g_{00} = c^2 \left(1 - \frac{2GM}{rc^2} \right)$$



$$\Rightarrow ds = \sqrt{c^2 \left(1 - \frac{2GM}{rc^2} \right)} dt \quad \Rightarrow \quad dt \simeq \frac{ds}{c} \left(1 + \frac{GM}{rc^2} \right)$$

- GR Length Contraction:

$$g_{ij} = -\delta_{ij} \left(1 + \frac{2GM}{rc^2} \right)$$

$$\Rightarrow ds = \sqrt{- \left(1 + \frac{2GM}{rc^2} \right) dx^i dx^i} \quad \Rightarrow \quad dt \simeq ds \left(1 - \frac{GM}{rc^2} \right)$$

The GR Deflection Angle, α_{GR}

- Using the metric $g_{\mu\nu}$ we can derive the Christoffel symbols using

$$\Gamma_{\mu\nu}^{\beta} = \frac{g^{\beta\beta}}{2} \left[\frac{\partial g_{\beta\mu}}{\partial x^{\nu}} + \frac{\partial g_{\beta\nu}}{\partial x^{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x^{\beta}} \right]$$

- This enables us to express the geodesic equation in terms of the metric:

$$\begin{aligned} \frac{E^2}{c^2} \frac{d^2 x^i}{dt^2} &= -\Gamma_{\mu\nu}^i p^{\mu} p^{\nu} \\ &= -\Gamma_{00}^i p^0 p^0 - 2\Gamma_{0j}^i p^0 p^j - \Gamma_{jk}^i p^j p^k \\ &= -\frac{MGx^i}{r^3} \frac{E^2}{c^2} - 0 - \Gamma_{33}^i p^3 p^3 \\ &= -\frac{MGx^i}{r^3} \frac{E^2}{c^2} - \frac{MGx^i}{r^3} \frac{E^2}{c^2} \\ &= -\frac{2MGx^i}{r^3} \frac{E^2}{c^2} \end{aligned}$$

Central term 2nd order
Last term only solution in z
using $p^z = E/c$ for photon

The GR Deflection Angle, α_{GR}

General Relativity

$$\frac{d^2 x^i}{dt^2} = -\frac{2MGx^i}{r^3}$$


Newtonian Gravity

$$\frac{d^2 \vec{r}}{dt^2} = -\frac{MG}{r^2} \vec{r}$$

- Hence, by realizing that x^i just represent the spatial vector of the photon in GR, we can move on from here, following the Newtonian derivation of α_{N} step by step carrying through the factor of 2 and eventual arrive at:

$$\alpha_{\text{GR}} = 2 \times \alpha_{\text{N}} = \frac{4MG}{\xi c^2}$$


So in summary...



$$\alpha_N = \frac{2MG}{\xi c^2}$$


$$\alpha_{GR} = 2 \times \alpha_N = \frac{4MG}{\xi c^2}$$

As claimed last week...




PHY-765 SS18 Gravitational Lensing Week 2

Questions?



PHY-765 SS18 Gravitational Lensing Week 2

Last Week's Worksheet



PHY-765 SS18 Gravitational Lensing Week 2

This Week's Worksheet