

PHY-765 SS18 Gravitational Lensing Week 4

Multiple Images

Kasper B. Schmidt

Leibniz-Institut für Astrophysik Potsdam (AIP)

Last week

• We derived the lens equation:

$$oldsymbol{eta} = oldsymbol{ heta} - oldsymbol{lpha}(oldsymbol{ heta})$$

• A source with true position β on the sky can be seen by an observer to be located at angular position θ under the deflection $\alpha(\theta)$.

• And defined (for the point mass) the

Critical Mass Surface Density Convergence

 $\Sigma_{\rm cr} \equiv \frac{c^2}{4\pi G} \frac{D_{\rm S}}{D_{\rm L} D_{\rm LS}} \qquad \kappa(\theta) \equiv \frac{\Sigma(D_{\rm L}\theta)}{\Sigma_{\rm cr}}$

Einstein Radius

$$heta_E \equiv \sqrt{rac{4MG}{c^2}}rac{D_{
m LS}}{D_{
m S}D_{
m L}}$$

The aim of today

- Explore the first consequence of the lens equation: multiple images
- Describe this for a few simplistic lens models
- Introduce the concepts of critical curves and caustics
- SN Refsdal a spectacular example of multiple images

Multiple Images from the Point Mass Lens

• Last week we described the point mass lens:

$$heta_E \equiv \sqrt{rac{4MG}{c^2}} rac{D_{
m LS}}{D_{
m S} D_{
m L}} \qquad oldsymbol{lpha}(oldsymbol{ heta}) = rac{4MG}{c^2} rac{D_{
m LS}}{D_{
m S} D_{
m L}} rac{oldsymbol{ heta}}{|oldsymbol{ heta}|^2} = rac{ heta_{
m E}^2}{|oldsymbol{ heta}|^2} oldsymbol{ heta}$$

• So we can write the lens equation as:

$$oldsymbol{eta} = oldsymbol{ heta} - rac{ heta_{
m E}^2}{|oldsymbol{ heta}|^2}oldsymbol{ heta}$$

_



- The lens equation is solved for all $\theta = \theta_{\rm E}$
- If imperfect alignment then x and y components of the lens equation are:

$$\beta = \theta_x \left[1 - \frac{\theta_{\rm E}^2}{\theta^2} \right] \qquad \qquad 0 = \theta_y \left[1 - \frac{\theta_{\rm E}^2}{\theta^2} \right]$$

- Assuming coordinate system aligned such that $\beta = \beta \hat{x}$ and $\beta > 0$



Multiple Images from the Point Mass Lens

- If $\theta_y \neq 0$, then we would have $\theta^2 \equiv \theta_x^2 + \theta_y^2 = \theta_E^2$
- But then $\beta = 0$ is violating $\beta > 0$
- $\beta = \theta_x \left[1 \frac{\theta_{\rm E}^2}{\theta^2} \right]$ So we must conclude that $\theta_v = 0$ (for the simple point mass lens)
 - I.e. the lens equation mapping is determined solely by the x-component



 $0 = \theta_y \left| 1 - \frac{\theta_{\rm E}^2}{\theta^2} \right|$

Multiple Images from the Point Mass Lens

• The limits for this setup are therefore:

$$egin{aligned} & heta_\pm \simeq \pm heta_{
m E} + rac{eta}{2} & (eta \ll heta_{
m E}) \ & heta_\pm = rac{eta}{2} \left[1 \pm \sqrt{1 + rac{4 heta_{
m E}^2}{eta^2}}
ight] \ & heta_\pm \simeq eta + rac{eta_{
m E}}{eta} & \& heta_- \simeq -rac{eta_{
m E}^2}{eta} & (eta \gg heta_{
m E}) \end{aligned}$$

• Using the Taylor expansion $\sqrt{1+\epsilon} \simeq 1 + \frac{1}{2}\epsilon - \dots$ for the limit $\beta \gg \theta_{\rm E}$

Spherically Symmetric Mass Distribution

- To start generalizing these ideas, we first look at the spherical distribution
- For a spherical distribution the convergence is independent on the direction

$$\kappa(\boldsymbol{\theta}) \equiv \frac{\Sigma(D_{\rm L}\boldsymbol{\theta})}{\Sigma_{\rm cr}} \longrightarrow \kappa(\boldsymbol{\theta}) \equiv \frac{\Sigma(D_{\rm L}\boldsymbol{\theta})}{\Sigma_{\rm cr}}$$

• Such that the deflection angle becomes

$$\boldsymbol{\alpha}(\boldsymbol{\theta}) = \frac{1}{\pi} \int d^2 \theta' \kappa(\theta') \frac{\boldsymbol{\theta} - \boldsymbol{\theta}'}{|\boldsymbol{\theta} - \boldsymbol{\theta}'|^2}$$

• $\alpha(\theta)$ is a vector. Only relevant vector is θ (κ doesn't care) so we can write:

$$\boldsymbol{\alpha}(\boldsymbol{\theta}) = A(\theta) \, \boldsymbol{\theta}$$

- $A(\theta)$ can be determined considering
 - The divergence on α :
 - Gauss' (divergence) theorem in the plane:

$$oldsymbol{
abla} oldsymbol{
abla} \cdot oldsymbol{lpha}(oldsymbol{ heta}) \ \int_{S} d^{2} heta \nabla \cdot oldsymbol{lpha}(oldsymbol{ heta}) = \oint_{\partial S} oldsymbol{ heta} \cdot oldsymbol{lpha}$$

J

Spherically Symmetric Mass Distribution

- This gives that $\nabla \cdot \boldsymbol{\alpha} = 2\kappa(\theta)$ $A(\theta) = \langle \kappa(\theta) \rangle$
- Where the mean normalized surface density has been defined as:

$$\langle \kappa(\theta)
angle = rac{1}{\pi \theta^2} \int_{\theta'- heta} d^2 \theta' \kappa(\theta')$$

• So we can express the lens equation for the spherical symmetric mass as

$$oldsymbol{eta} = oldsymbol{ heta} - \langle \kappa(heta)
angle oldsymbol{ heta}$$

• As κ is the ratio between surface density at angular distance θ from the lens (normalized by the critical surface density) this dictates that:

The deflection ($\beta - \theta$) a distance θ from the lens is governed by the mass contained within the cylinder of radius $\xi = D_L \theta$.

• which gives that

$$\langle \kappa(\theta)
angle = rac{M(R = D_{\rm L}\theta)}{\pi D_{\rm L}^2 \theta^2 \sum_{\rm cr}}$$

The Isothermal Sphere (IS)

• The result of the spherical symmetric mass distribution can be applied to

$$ho(r) = rac{\sigma^2}{2\pi G(r^2 + r_{
m core}^2)}$$

- r_{core} is the core radius (the density profile turns flat in the core)
- σ is the velocity dispersion of the lens



K.B. Schmidt, kbschmidt@aip.de

The Isothermal Sphere (IS)

• Using this density profile we have the surface density:

$$\Sigma(R) = \frac{\sigma^2}{2\pi G} \int_{-\infty}^{+\infty} \frac{dz}{R^2 + z^2 + r_{\rm core}^2} \quad \longrightarrow \quad \Sigma(R) = \frac{\sigma^2}{2G\sqrt{R^2 + r_{\rm core}^2}}$$

• Which can be used to express the average surface density within a radius R

$$M(R) = 2\pi \int_0^R dR' R' \Sigma(R') \quad \rightarrow \quad M(R) = \frac{\pi \sigma^2}{G} \left[\sqrt{R^2 + r_{\rm core}^2} - r_{\rm core} \right]$$

• Using $\theta = R/D_L$ and $\theta_{core} \equiv r_{core}/D_L$ we get the lens equation

$$\boldsymbol{\beta} = \boldsymbol{\theta} - \frac{\theta_0}{\theta^2} \left[\sqrt{\theta^2 + \theta_{\text{core}}^2} - \theta_{\text{core}} \right] \boldsymbol{\theta}$$

- by defining:

$$\theta_0 \equiv \frac{4\pi\sigma^2 D_{\rm LS}}{D_{\rm S}c^2}$$

The Singular Isothermal Sphere (SIS)

• For an isothermal sphere with no core ($\theta_{core} = 0$) the lens equation becomes

$$\boldsymbol{\beta} = \boldsymbol{\theta} \left[1 - \frac{\theta_0}{|\theta|} \right]$$

- where $\beta = 0$ generates an image (Einstein) ring motivating the θ_0 definition



Cored Isothermal Sphere (CIS)

- The mass density of real galaxies does not rise all the way into the center
- So even though SIS is simple, the assumption that $r_{core} = 0$ is poor.
- Using the definition of θ_0 and solving the lens equation for $\beta = 0$ we have

$$\theta^2 = \theta_0 \left[\sqrt{\theta^2 + \theta_{\rm core}^2} - \theta_{\rm core} \right] \qquad \qquad \theta_{\rm E} = \theta_0 \sqrt{1 - 2 \frac{\theta_{\rm core}}{\theta_0}}$$

- Hence, the size of the core determines when an Einstein ring can exist
 - I.e., if $2\theta_{core} > \theta_0$ then an Einstein ring cannot be formed
- Happens at ~1" for lens at 1Gpc
 - Actual physical size differes as θ_0 depends on z_{L} , z_{S} and σ
- If the lens-source alignment is not perfect the lens equation becomes

$$heta(eta- heta)=- heta_0\left[\sqrt{ heta^2+ heta_{
m core}^2}- heta_{
m core}
ight]$$

- Rather complex to solve for individual images.

Cored Isothermal Sphere (CIS)



• In 1D this special value $\beta_{caustic}$ is on a curve, but more generally it defines a circle of radius $\beta_{caustic}$ in the source plane

Caustics

- We want to determine the source position $\beta_{caustic}$
- Returning to the lens eq. on the form $\boldsymbol{\beta} = \boldsymbol{\theta} \frac{\theta_0}{\theta^2} \left[\sqrt{\theta^2 + \theta_{\text{core}}^2} \theta_{\text{core}} \right] \boldsymbol{\theta}$
- $\beta \rightarrow \pm \infty$ for $\theta \rightarrow \pm \infty$ so for three solutions there must be 2 extrema
- All values of β below $\beta(\theta_{-}) \equiv \beta_{\text{caustic}}$ produces 3 images (where $\beta > 0$)



Critical Curves



- The critical curves are define as the curves in the *lens* (image) plane where the images fall if the source is on the caustic in the source plane.
- Will return to caustic and critical curves when talking about magnification...

SN Refsdal

• Nov 2014: Discovered in MACS1149 data from the GLASS program





Existing Imaging

GLASS F104W

- Dec 2016: Imaging and spectroscopic follow-up of MACS1149 FoV
 - HFF, MOSFIRE, X-SHOOTER, DEIMOS, WFC3-G141

SN Refsdal



SN Refsdal - Multiple images

Kelly+2015



So in summary...

• The multiple images occurring from a point mass lens are given by

$$\theta_{\pm} = rac{\beta}{2} \left[1 \pm \sqrt{1 + rac{4\theta_{\mathrm{E}}^2}{\beta^2}} \right]$$

• The lens equation for the more general 'spherical lens' is

$$\boldsymbol{\beta} = \boldsymbol{\theta} - \langle \kappa(\theta) \rangle \boldsymbol{\theta} \quad \text{where} \quad \langle \kappa(\theta) \rangle = \frac{1}{\pi \theta^2} \int_{\theta' - \theta} d^2 \theta' \kappa(\theta') = \frac{M(R = D_{\rm L}\theta)}{\pi D_{\rm L}^2 \theta^2 \sum_{\rm cr}}$$

• This leads to the lens equations for the CIS and SIS:

$$\boldsymbol{\theta} = \boldsymbol{\theta} - \frac{\theta_0}{\theta^2} \left[\sqrt{\theta^2 + \theta_{\text{core}}^2} - \theta_{\text{core}} \right] \boldsymbol{\theta} \qquad \qquad \boldsymbol{\beta} = \boldsymbol{\theta} \left[1 - \frac{\theta_0}{|\theta|} \right]$$

- Solving for $\beta = 0$ reveals the multiple images for these lens models
- *Caustics* and *Critical curves* describe the source and image positions of multiple-image geometries in the source and lens (image) planes, respectively.
- SN Refsdal: a spectacular case of multiple images on galaxy and lens scales.



PHY-765 SS18 Gravitational Lensing Week 4

Questions?